



Danbury Hospital

Horblit Health Sciences Library

Warning Concerning Copyright Restrictions

No further reproduction or distribution of this copy is permitted by electronic transmission or any other means.

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material.

Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction not be "used for any purposes other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

This institution reserves the right to refuse to accept a copying order if, in its judgement, fulfillment of the order would involve violation of copyright law.

24475724

Request # 24475724**MAR 13, 2008****Email (PDF) To: NMCP-libraryservices@med.navy.mil**

US Naval Medical Center

Health Sciences Library Interlibrary Loan

620 John Paul Jones Circle

Portsmouth, VA 23708-2197

DOCLINE: Journal Copy

Title: Surgery for obesity and related diseases : official journal of the American Society for Bariatric Su
Title Abbrev: Surg Obes Relat Dis
Citation: 2008 Jan 31; [Epub ahead of print]
Article: Studies of Swedish adjustable gastric band and Lap
Author: Cunneen SA;Phillips E;Fielding G;Banelb S D;Estok R;Fahrbach
NLM Unique ID: 101233161 Verify: PubMed
PubMed UI: 18243061
ISSN: 1550-7289 (Print)
Publisher: Elsevier,, New York, N.Y. :
Copyright: Copyright Compliance Guidelines
Authorization: Lisa Eblen
Need By: N/A
Maximum Cost: **\$12.00**
Patron Name: Chastanet, [71]
Referral Reason: Not owned (title)
Library Groups: DOD,NAVY,THSL,FreeShare
Phone: 1.757.953-5386
Fax: 1.757.953-7533
Email: NMCP-libraryservices@med.navy.mil
Alternate Delivery: Email(PDF),Email(TIFF),Fax,Mail,Web(PDF)
Comments: **Thanks! #789**
Routing Reason: Routed to CTUDBH in Serial Routing - cell 5
Received: Mar 20, 2008 (11:05 AM EST)
Lender: Danbury Hospital/ Danbury/ CT USA (CTUDBH)

This material may be protected by copyright law (TITLE 17,U.S. CODE)

Bill to: VAUNRP

US Naval Medical Center

Health Sciences Library Interlibrary Loan

620 John Paul Jones Circle

Portsmouth, VA 23708-2197



Review

Studies of Swedish adjustable gastric band and Lap-Band: systematic review and meta-analysis

Scott A. Cunneen, M.D.^{a,*}, Edward Phillips, M.D.^a, George Fielding, M.D.^b,
Deirdre Banel, B.S.^c, Rhonda Estok, R.N., B.S.N.^c, Kyle Fahrbach, Ph.D.^c,
Isabella Sledge, M.D., M.P.H.^c

^aCedars Sinai Medical Center, Los Angeles, California;

^bDepartment of Surgery, New York University Medical Center, New York, New York;

^cUnited BioSource Corporation, Medford, Massachusetts

Received July 16, 2007; revised October 7, 2007; accepted October 19, 2007

Abstract

Background: This is the first systematic review and meta-analysis of the large body of data describing the Swedish adjustable gastric band (SAGB) and Lap-Band (LB).

Methods: A systematic review was performed that included screening of studies published in any language (January 1, 1998 through April 30, 2006) identified through MEDLINE, Current Contents, or the Cochrane Library. Studies with ≥ 10 SAGB or LB patients reporting ≥ 30 -day efficacy or safety outcomes were eligible for review; the data were extracted from the accepted studies. A weighted means analysis and random-effects meta-analysis of efficacy outcomes of interest were conducted.

Results: A total of 4592 bariatric surgery studies met the initial criteria. Of these studies, 129 (28,980 patients) were accepted (33 SAGB and 104 LB studies); most had a retrospective single-center design. For 4273 patients (36 treatment groups) in 33 SAGB studies and 24,707 patients (111 groups) in 104 LB studies, the mean baseline age (39.1–40.2 yr), body mass index (43.8–45.3 kg/m²), and gender (women 79.2–82.5%) were similar. A laparoscopic technique was used in $\geq 88\%$ and a pars flaccida technique in $\geq 41\%$ of both groups. Early mortality was equivalent for SAGB/LB ($\leq 1\%$). The 3-year mean SAGB and LB excess weight loss (56.36% and 50.20%, respectively) and body mass index reduction (-11.99 and -11.81 kg/m², respectively) from baseline were statistically significant ($P < .05$), as was the resolution of diabetes (61.45% and 60.29%, respectively) and hypertension (62.95% and 43.58%, respectively). Although scant and inconsistently reported data precluded direct statistical comparisons, the complication rates for the 2 devices appeared comparable. In 8 directly comparative studies, meta-analysis found a significantly greater absolute weight loss ($P < .05$) with the SAGB at 2 years (48.4 versus 41.9 kg, mean difference -4.84 , 95% confidence interval -9.47 to -0.22), although no difference was found in the percentage of excess weight loss or change in body mass index.

Conclusion: In a systematic review of the published world SAGB and LB data, at 1, 2, and 3 years, the weight loss, resolution of diabetes and hypertension, and complications appeared comparable. (Surg Obes Relat Dis 2008;xx:xxx–xxx.) © 2008 American Society for Metabolic and Bariatric Surgery. All rights reserved.

Keywords:

Morbid obesity; Laparoscopic adjustable gastric band; LAGB; Swedish adjustable gastric band; Lap-Band; Weight; Systematic review; Meta-analysis

This work was supported by Ethicon Endo-Surgery, a Johnson & Johnson Company, Cincinnati, Ohio

Presented at the 24th Annual Meeting of the American Society for Bariatric Surgery, June 11–16, 2007, San Diego, California

*Reprint requests: Scott A. Cunneen, M.D., Center for Weight Loss,

Cedars Sinai Medical Center, 8635 West Third Street, Suite 795W, Los Angeles, CA 90048.

E-mail: scott.cunneen@cshs.org

At the outset of the 21st century, bariatric surgery is the best available therapy [1–5] for the global epidemic of morbid obesity (body mass index [BMI] ≥ 40 or ≥ 35 kg/m² in the presence of high-risk co-morbidities) [6–9]. Implanted in >100,000 procedures [10], laparoscopic adjustable gastric banding (LAGB), a restrictive operation [11], is 1 of several weight-loss operations (i.e., LAGB, vertical banded gastroplasty, gastric bypass, biliopancreatic diversion/duodenal switch) to have demonstrated efficacy characterized by clinically meaningful ($\geq 50\%$) excess weight loss and co-morbidity reduction in the short (1–3 yr) and intermediate (≥ 3 –8 yr) terms [12–17]. LAGB also carries the benefits of minimal invasiveness [18], is associated with a premier perioperative safety record [12,19–23], is frequently selected by patients [24], and has the lowest mortality ($< .05\%$) of the bariatric procedures [14,21,22].

LAGB patients benefit from swift operative times of 30–60 minutes [25], minimal postoperative pain, a brief hospital stay, and a relatively rapid recovery involving minimal scarring. The hallmark of the LAGB procedure is an adjustability that facilitates individualized, modifiable, external band restriction such that short-term changes in nutritional requirements (e.g., pregnancy, illness) can be accommodated [13,26,27]. The LAGB procedure is also reversible, allowing restoration of essentially normal gastrointestinal anatomy and function or conversion to an alternate bariatric procedure [28–32]. The long-term outcomes (> 8 –10 yr) of LAGB, similar to those of other bariatric procedures, remain to be demonstrated [15].

Of the several gastric bands available globally, the 2 most often implanted are the Swedish adjustable gastric band (SAGB, Obtech Medical Sarl, Le Locle, Switzerland and Ethicon Endo-Surgery, Cincinnati, OH) and the Lap-Band (LB; Allergan Medical, Irvine, CA). The SAGB and adjustable gastric banding were introduced in the mid-1980s by Hallberg and Forsell [33] and Kuzmak [34], respectively, and have been consistently reported on in medical publications since then. The LB and SAGB were first implanted laparoscopically in 1993, reported on in 1995 by Belachew et al. [35] and Catona et al. [36] and in 1996 by Forsell and Hellers [37] and Klaiber et al. [38], and are currently the most used and widely studied bands. The SAGB is an investigational device in the United States as of July 7, 2007. The present study sought to assess the efficacy and safety of LAGB as a therapeutic strategy and to compare the SAGB and LB in terms of weight loss, co-morbidity improvement, and complications using a systematic review and meta-analysis of the world data.

Methods

Sources

In July 7, 2003, a prospective protocol was developed to undertake a systematic review [39] of the published medical

studies to identify and meta-analytically compare the SAGB and LB outcomes [40]. A broad electronic search was performed using PubMed accessing MEDLINE, Current Contents, and the Cochrane Library, with cutoff dates of January 1, 1998 through April 30, 2006. The following search terms were used: Obesity/surgery [MeSH] OR gastroplasty OR bariatric OR gastric banding, with the delimitation: human, 1998–2006. Manual supplementary reference checks of the accepted studies and reviews within the past 2 years were also performed.

Screening

Two levels of screening were applied (i.e., abstracts for all retrieved studies and full-text review for accepted studies). The acceptance requirements included studies of any design on ≥ 10 patients who had undergone SAGB or LB implantation and reporting extractable outcomes of interest. The bands must have been placed laparoscopically in $\geq 85\%$ of patients. The efficacy outcomes (i.e., percentage of excess weight loss, change in BMI, and reduction in absolute weight) were extracted at the 1-, 2-, 3-, and 5-year points, and at the latest point at which $\geq 50\%$ of the population had follow-up data available. Both early and late safety data (i.e., mortality and band/port/other complications) were required to be extractable by band type. Kin study data (multiple publications using the same study population) were counted only once, as part of the accepted study tabulation. All accepted studies were identified with a Centre for Evidence-Based Medicine (Oxford, UK) level of evidence from I to IV, with I being the best/most complete.

Statistical analysis

The efficacy outcomes of interest were synthesized using meta-analytic pooling of the SAGB and LB results. Meta-analyses of all efficacy outcomes were conducted using a random-effects model. The efficacy outcomes included both raw mean before and after changes (e.g., absolute weight loss) and proportions (e.g., response rates) [41–43]. Bayesian meta-analyses were also conducted for select outcomes to ascertain the robustness of the results. Heterogeneity was assessed using Cochrane's Q statistic. Zeroes were not assumed when not mentioned; rather, the data were treated as missing and were not included in the analyses. Zeroes were imputed for mortality calculations if all patients had been followed up and accounted for.

For studies in which SAGB and LB were studied side-by-side and directly compared, a meta-analysis was conducted for the mean before and after change differences (for continuous outcomes) and odds ratios (for binary outcomes). A positive difference in the before and after changes denoted a greater decrease in a continuous outcome measure in question. An odds ratio of < 1 denoted that a given binary outcome occurred more frequently. The meta-analytic mean values and weighted mean differences with

Table 1
Study characteristics

Study characteristic	Total			SAGB			Lap-Band			Direct comparison studies		
	k	t	n	k	t	n	k	t	n	k	t	n
Total	129	147	28,980	33	36	4,273	104	111	24,707	8	17	1,696
Study location												
Europe	83	95	18,416	25	27	3,601	64	68	14,815	6	13	1,304
North America	23	25	5,039	—	—	—	23	25	5,039	—	—	—
Other*	23	27	5,525	8	9	672	17	18	4,853	2	4	392
Study design												
RCT	8	12	478	4	6	209	5	6	269	1	2	180
Nonrandomized comparative study	27	35	3,626	10	11	748	22	24	2,878	5	10	528
Single-arm study	93	99	24,832	19	19	3,316	76	80	21,516	2	5	988
Case control	1	1	44	—	—	—	1	1	44	—	—	—
Prospective	46	55	6,640	14	17	1,447	35	38	5,193	3	7	1,040
Retrospective	83	92	22,340	19	19	2,826	69	73	19,514	5	10	656
Evidence level†												
I	5	8	520	2	3	130	4	5	390	1	2	180
II	16	19	1,108	7	9	489	10	10	619	1	2	156
III	31	36	4,665	7	7	400	27	29	4,265	3	6	320
IV	77	84	22,687	17	17	3,254	63	67	19,433	3	7	1,040
Institutional setting												
Single	114	130	21,801	30	33	3,827	91	97	17,974	7	15	1,516
Multicenter	15	17	7,179	3	3	446	13	14	6,733	1	2	180
Mean follow-up (mo)												
<24	44	51	4,645	12	13	1,002	34	38	3,643	2	5	988
24–35	36	40	6,280	10	11	1,165	29	29	5,115	3	6	365
>36	36	41	14,687	9	10	2,045	29	31	12,642	2	4	291
NR	13	15	3,368	2	2	61	12	13	3,307	1	2	52

SAGB = Swedish adjustable gastric band; k = number of studies; t = number of treatment groups; n = number of patients; RCT = randomized controlled trial; NR = not reported.

* Australia, Egypt, Israel, Saudi Arabia, and Singapore.

† Level of evidence according to Center of Evidence Based Medicine.

95% confidence intervals (CIs), were computed. For calculating the weight loss and co-morbidity outcomes, the latest point at which $\geq 50\%$ patient follow-up data were available was used. Early (<30 d) and late (≥ 30 d) complications were summarized by band type and whether the pars flaccida technique was used. Calculations were performed using Statistical Analysis Systems software, version 8.1 (SAS Institute, Cary, NC) and Statistical Package for Social Sciences, version 13.0 (SPSS, Chicago, IL).

Results

Screening outcomes

Of the complete catalogue of SAGB and LB citations identified ($n = 4594$), level 1 screening rejected 2740 studies. Of the remaining 1854 studies, 1549 were rejected in the level 2 screening. Of the 305 remaining studies, 176 were kin studies for which the data were counted only once in amalgamation with the overlapping data of the 129 “accepted studies” (28,980 patients), all of which were fully extracted and eligible for meta-analysis (see the Appendix of Accepted Studies).

Study characteristics

The accepted studies included 83 European studies (18,416 patients), 22 North American studies (5039 patients), and 24 studies from other locations (Australia, Egypt, Israel, Saudi Arabia, Singapore; Table 1). Most studies were retrospective case series conducted at single centers. Of the studies reporting a median (or mean) follow-up time, 44 reported <2 years of follow-up, 36 studies reported 2–3 years, 36 reported >3 years of follow-up, and 13 did not report on the length of follow-up. SAGB was represented in 36 treatment groups (4273 patients) in 33 studies and LB in 111 treatment groups (24,707 patients) in 104 studies.

Patient characteristics

The baseline demographics were comparable. The mean age, percentage of women, and mean BMI was 39.1 and 40.2 years, 79.2% and 82.5%, and 43.8 and 45.3 kg/m^2 for the SAGB and LB groups). Most patients met the National Institutes of Health [6] criteria for weight loss surgery. The percentages of patients with co-morbidities at baseline were more varied (e.g., 24% of 1083 SAGB patients had hyper-

tension in the 8 treatment arms reporting this characteristic compared with 37% of 3664 LB patients in 28 treatment arms). Too few treatment arms reported on the SAGB baseline co-morbidities to draw meaningful conclusions about patient differences between the groups. In studies providing baseline co-morbidity data, rarely was improvement or resolution reported at follow-up. Also, the follow-up duration varied widely (range 1–98 mo).

A subset of 8 studies (1696 patients) directly compared SAGB and LB [44–51], of which only 1 was a randomized comparison [51]. Of the 6 European studies and 2 from other countries, 3 presented their 3-year results. The SAGB/LB patient baseline demographics were similar to those of the primary accepted studies.

Technique and hospitalization

The vast majority of SAGB and LB patients underwent laparoscopic band placement (87.9% and 98.8%, respectively), with a similar (1.1% and 2.0%) rate of conversion to an open procedure. The pars flaccida technique was used in 16 of 36 SAGB and 28 of 111 LB groups. The investigators often did not explicitly state the technique used, and the accrual years for many cohorts were broad and included both the pars flaccida and the perigastric techniques. The mean length of hospital stay was not significantly different between the groups.

Weight loss

Significant weight reduction ($P < .05$) was observed in the SAGB and LB groups independently when computations using either simple weighted pooling of the estimates of all studies or meta-analytic computations of the mean change were performed (Table 2). At a follow-up of 1, 2, and 3 years, the bands demonstrated steady weight loss, reaching 56.36% (95% CI 48.14–64.58%) and 50.20% (95% CI 43.36–57.0%) excess weight loss in the SAGB and LB groups at 3 years, respectively. In the subset of comparative studies, a meta-analytic comparison at 2 years and at the last point at which $\geq 50\%$ of patients had follow-up data available showed a statistically significant ($P < .05$) mean difference in absolute weight loss (weighted mean: SAGB, -48.38 versus LB, -41.9 ; weighted mean difference: -4.84 , 95% CI -9.47 to -0.22); at the last point reported, however, the weight loss measures varied widely, from 4 to 36 months.

The change in BMI was consistent and statistically significant ($P < .05$) from the baseline value for both SAGB and LB, independently, at all points (3 yr -11.99 , 95% CI -15.20 to -8.77 and -11.81 , 95% CI -13.16 to -10.47 , respectively). These results were corroborated by a sensitivity analysis of only those SAGB studies with weight loss data at all points (data not shown). An upward bias of the weight loss results because of selective dropout of patients with weight loss failure did not appear to contribute to the

findings; the numbers of patients available at each point for the sensitivity analysis were not substantively reduced. Too few studies and heterogeneity of the SAGB and LB percentage of excess weight loss and BMI data precluded meta-analytic comparison of these variables. The weight loss results from the meta-analyses using Bayesian method was not substantively different from the results using a random effects meta-analysis.

Co-morbid disease

For the specific co-morbidities of obesity and their response to LAGB, the number of studies available for either band was somewhat low and the definitions used of changes in categorical outcomes (e.g., resolved, improved, resolved/improved, unchanged, worsened) were often inconsistent, unclear, or not stated. The time frames for improvement were also inconsistently described; only the results for a change from baseline for type 2 diabetes and hypertension included a meaningful number of studies for reporting the meta-analysis (Table 3).

Two SAGB and 14 LB studies reported a statistically significant resolution of diabetes ($P < .05$) in 61.5% of 117 SAGB patients and 60.3% of 249 LB patients at 2 years for those who had diabetes at baseline. Significant reductions in insulin (-67.87 and -70.04 pmol/L, respectively) and fasting blood glucose (-0.89 and -0.63 mmol/L, respectively) provided laboratory substantiation of the clinical outcomes in the SAGB and LB studies. Patients also had high rates of hypertension resolution at 2 years—62.95% of patients in 5 SAGB treatment arms (228 patients) and 43.6% in 12 LB treatment arms (488 patients)—an improvement that persisted for >2 years in 42.9% of SAGB patients. No LB studies reported the results for hypertension after 2 years. The small number of studies precluded a meta-analytic comparison of the changes in co-morbidities.

Complications

Early mortality was equivalent for both devices (.1%), and late mortality was comparable (SAGB .2% versus LB .1%). The incidence of early and late complications was low. Overall, the complication rates for the SAGB and LB appeared similar (Table 4), although a statistical comparison of the complications was not possible owing to the widespread inconsistency in data reporting. For all studies describing complications, the frequency of late slippage or migration (4.0% and 6.2% for SAGB and LB, respectively) and pouch dilation (1.7% and 5.1% for SAGB and LB, respectively) was lower for SAGB. When the analysis controlled for the use of the pars flaccida technique, the differences in the complication rates were reduced (from 4.3% to 2.6% for SAGB and 6.9% to 3.1% for LB). Few other differences in SAGB and LB morbidity were apparent, and all were highly variable in the direction and magnitude of the difference. The number of studies reporting additional

Table 2
Total weight loss outcomes

Outcome	SAGB			Lap-Band		
	t (n)	Weighted mean	Mean change (95% CI)	t (n)	Weighted mean	Mean change (95% CI)
Absolute weight (kg)						
Last follow-up* (yr)	20 (1235)	−46.04	−40.35 (−48.91, −31.79)†	34 (2436)	−33.66	−30.14 (−34.54, −25.74)†
1	10 (877)	−39.22	−41.30 (−50.55, −32.05)†	26 (1541)	−29.12	−28.95 (−31.87, −26.03)†
2	9 (387)	−49.08	−44.98 (−57.18, −32.79)†	15 (803)	−33.21	−34.17 (−39.71, −28.64)†
3	4 (323)	−51.79	−49.89 (−64.18, −35.61)†	7 (501)	−33.93	−37.25 (−50.20, −24.31)†
5	—	—	—	2 (123)	−24.07	—
BMI (kg/m ²)						
Last follow-up*	26 (2459)	−12.25	−12.30 (−14.48, −10.13)†	61 (8217)	−11.05	−10.22 (−11.08, −9.37)†
1	16 (2778)	−9.89	−11.90 (−14.42, −9.38)†	51 (7804)	−10.75	−10.11 (−10.76, −9.46)†
2	13 (2164)	−11.51	−13.69 (−16.94, −10.44)†	37 (4275)	−11.67	−11.32 (−12.10, −10.54)†
3	6 (1560)	−11.10	−11.99 (−15.20, −8.77)†	25 (4495)	−13.00	−11.81 (−13.16, −10.47)†
5	2 (190)	−13.02	—	4 (249)	−11.32	−9.21 (−15.22, −3.20)†
%EWL						
Last follow-up*	12 (2221)	55.28	55.62 (50.04, 61.21)†	47 (6251)	43.90	43.57 (39.98, 47.15)†
1	9 (1943)	38.83	44.48 (37.35, 51.61)†	45 (6204)	42.44	42.97 (40.39, 45.56)†
2	6 (1657)	46.49	49.98 (45.86, 54.10)†	32 (3540)	50.38	48.14 (43.26, 53.01)†
3	6 (1560)	52.63	56.36 (48.14, 64.58)†	19 (1659)	54.13	50.20 (43.36, 57.04)†
5	2 (190)	56.71	—	6 (823)	51.14	39.76 (27.52, 52.00)†
Meta-analysis	Comparative studies only				WMD (95% CI)	
	t (n)	Weighted mean	t (n)	Weighted mean		
Absolute weight (kg)	3 (101)	−50.4	3 (147)	−44.07	−4.82 (−9.36, −0.27)‡	
Last follow-up*						
1	3 (127)	−41.32	3 (157)	−44.22	2.50 (−1.73, 6.74)	
2	3 (101)	−48.38	3 (134)	−41.9	−4.84 (−9.47, −0.22)‡	
3	2 (70)	−52.34	2 (112)	−45.96	—	
BMI (kg/m ²)						
Last follow-up*	2 (115)	−13.32	2 (123)	−11.1	—	
1	2 (130)	−9.76	2 (124)	−10.02	—	
2	2 (115)	−12.59	2 (109)	−13.12	—	
3	1 (84)	−13.40	1 (88)	−11.6	—	
%EWL						
Last follow-up*	2 (672)	51.88	3 (167)	51.24	—	
1	2 (677)	45.04	3 (168)	48.24	—	
2	1 (84)	55.00	1 (87)	63	—	
3	1 (84)	65.00	1 (88)	54	—	

SAGB = Swedish adjustable gastric band; t = number of treatment groups; n = number of patients evaluated; CI = confidence interval; BMI = body mass index; %EWL = percentage of excess weight loss; WMD = weighted mean difference (mean difference in pooled groups from comparison studies only).

* Last follow-up available with ≥50% of patients followed up.

† P < .001; significant mean effect.

‡ Significant mean effect.

band-related, port-related, or other complications was low, as were the rates of absolute events, rendering the clinical significance of these data uncertain.

In the subset of 8 studies reporting both SAGB and LB results, odds ratios were computed for specific complications, none of which reached statistical significance in favor of either band.

Discussion

This is the first systematic review of the world data (4595 citations) to encompass cumulative studies of the 2 most

commonly used LAGBs. The study designs varied and ranged from randomized controlled clinical trials to single-center retrospective reviews.

The most consistently and well-described data among the accepted studies were the weight loss measures. Both bands consistently evidenced statistically significant ($P < .05$) and durable, safe, weight loss (approximately 50% excess weight loss) at 3 years.

The resolution of diabetes and hypertension was significant for both devices and appeared to parallel weight loss at 2 years. As has been shown, the improvement and/or resolution of co-morbidities correlates with even

Table 3
Resolution of co-morbidities

Co-morbidity	SAGB		Lap-Band	
	t (n)	Mean (95% CI)	t (n)	Mean (95% CI)
Diabetes				
Resolved*	2 (117)	61.45 (52.11, 70.78)	14 (249)	60.29 (44.75, 75.83) ^{†‡}
Resolved or improved	—	—	13 (436)	83.24 (71.55, 94.93) ^{†‡}
Improved	—	—	13 (436)	35.99 (18.92, 53.06) ^{†‡}
Unchanged	—	—	9 (245)	16.37 (6.05, 26.69) ^{†‡}
Worsened	—	—	2 (54)	22.35 (0.00, 52.10) [§]
Diabetes measure				
Insulin (pmol/L)	4 (133)	−67.87 (−101.14, −34.60) [†]	4 (107)	−70.04 (−102.79, −37.28) ^{†‡}
HbA1c (%)	—	—	3 (99)	−0.37 (−0.83, 0.09) [§]
Fasting glucose (mmol/L)	5 (143)	−0.89 (−1.41, −0.37) [†]	3 (97)	−0.63 (−0.98, −0.29) [‡]
Hypertension				
Resolved*	5 (228)	62.95 (41.18, 84.72) [†]	12 (488)	43.58 (30.21, 56.95) ^{†‡}
Resolved or improved	1 (235)	74.89 (69.35, 80.44)	11 (942)	70.35 (55.26, 85.43) ^{†‡}
Improved	1 (235)	74.89 (69.35, 80.44)	11 (942)	41.36 (33.80, 48.91) ^{†‡}

SAGB = Swedish adjustable gastric band; t = number of treatment groups reporting characteristic; n = number of patients evaluated; CI = confidence interval; HbA1c = hemoglobin A1c.

* Discontinued treatment or, for diabetes improvement, reduced treatment.

[†] $P < .001$ for test of heterogeneity of outcome.

[‡] Statistically significant pre- vs. postoperative difference within surgery category.

[§] $P < .01$ for test of heterogeneity of outcome.

modest reductions (10–15%) of weight after bariatric surgery [17,52,53].

The common complications recorded for LAGB included gastric prolapse or slippage, pouch dilatation, erosion, and device breakage. The complication profiles for SAGB and LB appeared quite comparable. Very few serious early complications were associated with either band, in contrast to other bariatric operations. Both devices demonstrated minimal long-term complication and failure rates, as evidenced by similar durable weight loss.

In the 129 studies accepted for systematic review, the less well-known co-morbidities were either absent from the population or poorly documented. Future studies should clearly define the characteristics of change in co-morbid diseases measured and the time frames in which the data were recorded. Descriptions of postoperative management were limited, lacked standardization, and primarily comprised only the short or intermediate term. Only a few studies followed up patients for ≥ 6 –10 years and captured information on meaningful numbers of patients at those extended points [21,23,54–56]. Only 8 studies were identified that were designed to directly compare the SAGB and LB performances. Meta-analytic comparisons would be greatly strengthened by the availability of more SAGB studies and more comparative studies of SAGB and LB with intermediate-term (3–10 yr) and truly long-term (≥ 10 yr, as defined by O'Brien et al. [15]) follow-up for $\geq 50\%$ of patients.

Both SAGB and LB achieve weight loss and satiety by restriction. The SAGB has been, since its inception, engineered for “low-pressure” restriction, which could have a

bearing on the development of long-term complications, and requires additional study. Although not comparable statistically, the meta-analysis findings for both bands were in agreement with the most-cited banding data identifying LAGB, in general, as a safe bariatric procedure. The SAGB was originally designed for implantation using the pars flaccida pathway. This is a technique that has yielded low rates of long-term complications with the SAGB, and, as evidenced in this meta-analysis, has resulted in LB morbidity outcomes also significantly improved compared with those achieved using the perigastric approach [57].

The subset of 8 studies that directly compared SAGB and LB [44–51] included the prospective comparison of short-term SAGB/LB weight loss and complications by Ponson et al. [47], which found no statistically significant differences between these bands, and the prospective, randomized comparison of early SAGB/LB results by Suter et al. [51], which also found no difference in weight loss, food tolerance, or quality of life at 18 months and “identical” rates of late morbidity and reoperation. Each of the SAGB/LB comparative studies independently concluded band comparability.

The statistically significant findings in this meta-analysis for weight loss, resolution of co-morbidities, and mortality with respect to LAGB as a surgical approach to morbid obesity corroborate those of prominent systematic reviews of bariatric outcomes, such as Fielding et al. [12], O'Brien et al. [15], Fried et al. [58], and Chapman et al. [14]. Moreover, the outcomes for banding at 3 years in the present study (laparoscopic implantation for $\geq 88\%$; percentage of excess weight loss $\geq 50\%$; BMI reduction ≥ 11.81 kg/m²; diabetes and hypertension resolution of

Table 4
Complications

Complication	SAGB						Lap-Band					
	Total			Pars flaccida			Total			Pars flaccida		
	t	n/N	%	t	n/N	%	t	n/N	%	t	n/N	%
Mortality												
Early	24	4/3,825	0.1	12	4/1,990	0.2	66	15/14,518	0.1	21	4/5,585	0.1
Late	21	6/3,596	0.2	10	5/1,841	0.3	54	14/10,056	0.1	17	1/3,317	0.0
Band related												
Erosion												
Early	5	0/1,740	0.0	3	0/1,061	0.0	19	4/4,972	0.1	9	3/2,944	0.1
Late	10	23/2,141	1.1	5	20/1,386	1.4	30	140/10,840	1.3	10	42/3,500	1.2
Leakage												
Early	—	—	—	—	—	—	1	4/63	6.3	—	—	—
Late	1	2/625	0.3	—	—	—	5	6/415	1.4	2	1/237	0.4
Slippage/migration												
Early	10	8/2,301	0.3	4	4/1,411	0.3	35	66/8,945	0.7	10	41/2,800	1.5
Late	13	118/2,966	4.0	5	35/1,370	2.6	55	813/13,113	6.2	12	96/3,064	3.1
Port related												
Leakage												
Early	1	0/824	0.0	1	0/824	0.0	8	0/2,868	0.0	2	0/101	0.0
Late	2	19/914	2.1	2	19/914	2.1	16	87/4,331	2.0	5	18/824	2.2
Tubing rupture												
Early	1	0/34	0.0	1	0/34	0.0	9	35/3,954	0.9	6	35/2,404	1.5
Late	4	9/618	1.5	2	2/249	0.8	11	85/4,627	1.8	7	37/2,199	1.7
Dislocation												
Early	4	0/1,004	0.0	4	0/1,004	0.0	14	26/2,941	0.9	7	8/1,390	0.6
Late	10	69/2,781	2.5	6	20/1,359	1.5	27	150/4,920	3.0	7	14/935	1.5
Other												
Esophageal dilation												
Early	2	2/230	0.9	2	2/230	0.9	3	0/1,504	0.0	3	0/1,504	0.0
Late	3	11/1,026	1.1	1	2/90	2.2	11	91/5,583	1.6	4	16/1,754	0.9
Pouch dilation												
Early	8	7/863	0.8	4	4/215	1.9	22	51/3,002	1.7	3	0/224	0.0
Late	11	19/1,122	1.7	5	8/430	1.9	32	222/4,372	5.1	6	28/759	3.7
Stomal stenosis												
Early	—	—	—	—	—	—	11	57/3,687	1.5	4	47/2,924	1.6
Late	1	1/26	3.8	—	—	—	8	104/3,143	3.3	1	0/1,014	0.0

SAGB = Swedish adjustable gastric band; t = number of treatment groups; n = number of patients with outcome; N = number of patients evaluated; % = percentage of patients with complication.

≥60.29 and ≥43.58%, respectively; early mortality rate .1%) were clinically meaningful relative to the outcomes benchmarks established in medical studies, such as Buchwald et al. [17,59] and Maggard et al. [60], using the more stringent standard of meta-analysis. Although in this systematic review and meta-analysis of SAGB and LB, only absolute weight loss was appropriate for direct statistical comparison within a small subset of studies, the findings for LAGB as a device class are powerful, demonstrating its safety and statistically and clinically significant efficacy for the short to early-intermediate term.

Disclosures

The authors have no commercial associations that might be a conflict of interest in relation to this article.

Acknowledgment

We would like to acknowledge Jane Buchwald, M.A. and Roy Kulick, M.D. for their substantial contributions to the development of this report.

References

- [1] North American Association for the Study of Obesity and National Heart, Lung, and Blood Institute. *The practical guide: identification, evaluation, and treatment of overweight and obesity in adults. Publication No. 00-4084*. Bethesda: National Institutes of Health; 2000.
- [2] National Institutes of Health and National Heart, Lung, and Blood Institute. *Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: the evidence report. Publication No. 98-4083*. Bethesda: National Institutes of Health; 1998.

- [3] Pories WJ, Swanson MS, MacDonald KG, et al. Who would have thought it? An operation proves to be the most effective therapy for adult-onset diabetes mellitus. *Ann Surg* 1995;222:339–52.
- [4] Goodrick GK, Foreyt JP. Why treatments for obesity don't last. *JAMA* 1991;91:1243–7.
- [5] O'Brien PE, Dixon JB, Laurie C, et al. Treatment of mild to moderate obesity with laparoscopic adjustable gastric banding or an intensive medical program: a randomized trial. *Ann Intern Med* 2006;144:625–33.
- [6] NIH Consensus Development Conference Panel. NIH Consensus Development Conference Statement: gastrointestinal surgery for severe obesity. *Obes Surg* 1991;1:243–56.
- [7] Deitel M. Overweight and obesity worldwide now estimated to involve 1.7 billion people. *Obes Surg* 2003;13:329–30.
- [8] International Obesity Task Force. Call for obesity review as overweight numbers reach 1.7 billion (press release). London, England: International Obesity Task Force; March 17, 2003. Available from: <http://www.iotf.org/media/iotfmar17.htm>. Accessed September 12, 2004.
- [9] Buchwald H, Williams SE. Bariatric surgery worldwide 2003. *Obes Surg* 2004;14:1157–64.
- [10] Suter M, Calmes JM, Paroz A, Giusti V. A 10-year experience with laparoscopic gastric banding for morbid obesity: high long-term complication and failure rates. *Obes Surg* 2006;16:829–35.
- [11] Buchwald H, Buchwald JN. Evolution of operative procedures for the management of morbid obesity: 1950–2000. *Obes Surg* 2002;12:705–17.
- [12] Fielding GA, Ren CJ. Laparoscopic adjustable gastric band. *Surg Clin North Am* 2005;85:129–40.
- [13] O'Brien PE, Dixon JB, Brown W, et al. The laparoscopic adjustable gastric band (Lap-Band): a prospective study of medium-term effects on weight, health and quality of life. *Obes Surg* 2002;12:652–60.
- [14] Chapman AE, Kiroff G, Game P, et al. Laparoscopic adjustable gastric banding in the treatment of obesity: a systematic literature review. *Surgery* 2004;135:326–51.
- [15] O'Brien PE, McPhail T, Chaston TB, Dixon JB. Systematic review of medium-term weight loss after bariatric operations. *Obes Surg* 2006;16:1032–40.
- [16] Ponce J, Haynes B, Paynter S, et al. Effect of Lap-Band–induced weight loss on type 2 diabetes mellitus and hypertension. *Obes Surg* 2004;14:1335–42.
- [17] Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA* 2004;292:1724–37.
- [18] O'Connell T. An overview of obesity and weight-loss surgery. *Clin Diabetes* 2004;22:115–20.
- [19] Zinzindohoue F, Chevallier JM, Douard R, et al. Laparoscopic gastric banding: a minimally invasive surgical treatment for morbid obesity: prospective study of 500 consecutive patients. *Ann Surg* 2003;237:1–9.
- [20] Parikh MS, Laker S, Weiner M, Hajiseyedyavadi O, Ren CJ. Objective comparison of complications resulting from laparoscopic bariatric procedures. *J Am Coll Surg* 2006;202:252–61.
- [21] O'Brien PE, Dixon JB. Lap-Band: outcomes and results. *J Laparoendosc Adv Surg Tech* 2003;13:265–70.
- [22] Martikainen T, Pirinen E, Alhava E, et al. Long-term results, late complications and quality of life in a series of adjustable bands. *Obes Surg* 2004;14:648–54.
- [23] Weiner R, Blanco-Engert R, Weiner S, et al. Outcome after laparoscopic adjustable gastric banding—8 years experience. *Obes Surg* 2003;13:427–34.
- [24] Ren CJ, Cabrera I, Rajaram K, Fielding GA. Factors influencing patient choice for bariatric operation. *Obes Surg* 2005;15:202–6.
- [25] Mizrahi S, Avinoah E. Technical tips for laparoscopic gastric banding: 6 years' experience in 2800 procedures by a single surgical team. *Am J Surg* 2007;193:160–5.
- [26] Favretti F, O'Brien PE, Dixon JB. Patient management after Lap-Band placement. *Am J Surg* 2002;184:38S–41S.
- [27] Dixon JB, O'Brien PE. Nutritional outcomes of bariatric surgery. In: Buchwald H, Cowan SM, Pories WJ, editors. *Surgical management of obesity*. Philadelphia: WB Saunders; 2006. p. 357–64.
- [28] Westling A, Ohrvall M, Gustavsson S. Roux-en-Y gastric bypass after previous unsuccessful gastric restrictive surgery. *J Gastrointest Surg* 2002;6:206–11.
- [29] Weber M, Muller M, Michel JM. Laparoscopic Roux-en-Y gastric bypass, but not rebanding, as rescue procedure for patients with failed laparoscopic gastric banding. *Ann Surg* 2003;238:827–33.
- [30] Kothari SN, DeMaria EJ, Sugerman HJ. Laparoscopic band failure: conversion to gastric bypass and their preliminary outcomes. *Surgery* 2002;131:625–9.
- [31] Dolan K, Fielding GA. Biliopancreatic diversion following failure of laparoscopic adjustable gastric banding. *Surg Endosc* 2004;18:60–3.
- [32] Slater GH, Fielding GA. Combined laparoscopic adjustable gastric banding and BPD after failed bariatric surgery. *Obes Surg* 2004;14:677–82.
- [33] Hallberg D, Forsell P. Ballongband vid behandling av massiv övervikt (Balloon band for treatment of massive obesity). *Svensk Kirurgi* 1985;43:106.
- [34] Kuzmak LI. Silicone gastric banding: a simple and effective operation for morbid obesity. *Contemp Surg* 1986;28:13–8.
- [35] Belachew M, Legrand M, Vincent V, et al. Laparoscopic placement of adjustable silicone gastric band in the treatment of morbid obesity: how to do it. *Obes Surg* 1995;5:66–70.
- [36] Catona A, La Manna L, Forsell P. The Swedish adjustable gastric band: laparoscopic technique and preliminary results. *Obes Surg* 2000;10:15–21.
- [37] Forsell P, Hellers G. The Swedish adjustable gastric banding (SAGB) for morbid obesity: 9-year experience and a 4-year follow up of patients operated with a new adjustable band. *Obes Surg* 1997;7:345–51.
- [38] Klaiber C, Metzger A, Forsell P. Laparoscopic gastric banding. *Chirurg* 2000;71:146–51.
- [39] Cook DJ, Mulrow CD, Haynes RB. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med* 1997;126:376–80.
- [40] MetaWorks, Inc. Protocol: bariatric surgery using Swedish adjustable gastric band or Lap-Band: a systematic review of the literature and meta-analysis. July 2003.
- [41] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- [42] Hedges L, Olkin I. *Statistical methods for meta-analysis*. Orlando, FL: Academic Press; 1985. p. 230–57.
- [43] Al-Jiffry BO, Shaffer EA, Saccone GT, et al. Changes in gallbladder motility and gallstone formation following laparoscopic gastric banding for morbid obesity. *Can J Gastroenterol* 2003;17:169–74.
- [44] Miller K, Hell E. Laparoscopic adjustable gastric banding: a prospective 4-year follow-up study. *Obes Surg* 1999;9:183–7.
- [45] Nowara HA. Egyptian experience in laparoscopic adjustable gastric banding (technique, complications, and intermediate results). *Obes Surg* 2001;11:70–5.
- [46] Vassallo C, Andreoli M, La Manna A, Turpini C. 60 reoperations on 890 patients after gastric restrictive surgery. *Obes Surg* 2001;11:752–6.
- [47] Ponson AE, Janssen IM, Klinkenbijn JH. Laparoscopic adjustable gastric banding: a prospective comparison of two commonly used bands. *Obes Surg* 2002;12:579–82.
- [48] Bende J, Csiszar M, Ursu M. Modern surgical interventions in the treatment of morbid obesity. *Orv Hetil* 2003;144:2459–63.
- [49] Ceelen W, Walder J, Cardon A, et al. Surgical treatment of severe obesity with a low-pressure adjustable gastric band: experimental data and clinical results in 625 patients. *Ann Surg* 2003;237:10–6.

- [50] Dukhno O, Ovnat A, Levy I. Our experience with 250 laparoscopic adjustable silicone gastric bandings. *Surg Endosc* 2003;17:601–2.
- [51] Suter M, Giusti V, Worreth M, et al. Laparoscopic gastric banding: a prospective, randomized study comparing the Lap Band and the SAGB: early results. *Ann Surg* 2005;241:55–62.
- [52] U.S. Department of Health and Human Services. *The Surgeon General's call to action to prevent and decrease overweight and obesity. Section 1. Overweight and obesity as public health problems in America*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Office of the Surgeon General; 2001.
- [53] Christou NV, Sampalis JS, Lieberman M, et al. Surgery decreases long-term mortality, morbidity, and health care use in morbidly obese patients. *Ann Surg* 2004;240:416–23.
- [54] Fielding GA, Duncombe JE. Clinical and radiological follow-up of a cohort of laparoscopic adjustable gastric bands from 1998. *Obes Surg* 2005;15:634–40.
- [55] Vertruyen M. Experience with Lap-band system up to 7 years. *Obes Surg* 2002;12:569–72.
- [56] O'Brien PE, Dixon JB. Weight loss and early and late complications—the international experience. *Am J Surg* 2002;184:S42–S45.
- [57] Allen JW, Coleman MG, Fielding GA. Lessons learned from laparoscopic gastric banding for morbid obesity. *Am J Surg* 2001;182:10–4.
- [58] Fried M, Miller K, Kormanova K. Literature review of comparative studies of complications with Swedish band and Lap-Band. *Obes Surg* 2004;14:256–60.
- [59] Buchwald H, Estok R, Fahrbach K, Banel D, Sledge I. Trends in mortality in bariatric surgery: a systematic review and meta-analysis. *Surgery* 2007;142:621–35.
- [60] Maggard M, Shugarman LR, Suttrop M, et al. Meta-analysis: surgical treatment of obesity. *Ann Intern Med* 2005;142:547–59.
- [10] Belachew M, Legrand M, Vincent V, Lismonde M, Le Docte N, Deschamps V. Laparoscopic adjustable gastric banding. *World J Surg* 1998;22:955–63.
- [11] Bende J, Csiszar M, Ursu M. Modern surgical interventions in the treatment of morbid obesity. *Orv Hetil* 2003;144:2459–63.
- [12] Berg T, Zurmeyer EL, Ranneberg T, Schonleben K. Surgical therapy of morbid obesity using an adjustable gastric band: report of experiences over 2 1/2 years with 71 patients. *Med Klin (Munich)* 2001;96:191–5.
- [13] Blanco-Engert R, Weiner S, Pomhoff I, Matkowitz R, Weiner RA. Outcome after laparoscopic adjustable gastric banding, using the Lap-Band and the HelioGast band: a prospective randomized study. *Obes Surg* 2003;13:776–9.
- [14] Burhop JW, Chiang MC, Engstrand DJ, O'Driscoll M. Laparoscopic bariatric surgery can be performed safely in the community hospital setting. *WMJ* 2005;104:48–53.
- [15] Camerini G, Adami G, Marinari GM, et al. Thirteen years of follow-up in patients with adjustable silicone gastric banding for obesity: weight loss and constant rate of late specific complications. *Obes Surg* 2004;14:1343–8.
- [16] Capizzi FD, Boschi S, Brulatti M, et al. Laparoscopic adjustable esophagogastric banding: preliminary results. *Obes Surg* 2002;12:391–4.
- [17] Catona A, La Manna L, Forsell P. The Swedish adjustable gastric band: laparoscopic technique and preliminary results. *Obes Surg* 2000;10:15–21.
- [18] Ceelen W, Walder J, Cardon A, et al. Surgical treatment of severe obesity with a low-pressure adjustable gastric band: experimental data and clinical results in 625 patients. *Ann Surg* 2003;237:10–6.
- [19] Chau WY, Schmidt HJ, Kouli W, Davis D, Wasielewski A, Ballantyne GH. Patient characteristics impacting excess weight loss following laparoscopic adjustable gastric banding. *Obes Surg* 2005;15:346–50.
- [20] Chevallier JM, Zinzindohoue F, Douard R, Blanche JP, Berta JL, Altman JJ. Complications after laparoscopic adjustable gastric banding for morbid obesity: experience with 1,000 patients over 7 years. *Obes Surg* 2004;14:407–14.
- [21] Collet D, Rault A, Sa Cunha A, Larroude D, Masson B. Laparoscopic adjustable gastric banding results after 2 years with two different band types. *Obes Surg* 2005;15:853–7.
- [22] Coskun H, Bozboru A, Ogunc G, Peker Y. Adjustable gastric banding in a multicenter study in Turkey. *Obes Surg* 2003;13:294–6.
- [23] Coupaye M, Bouillot JL, Coussieu C, Guy-Grand B, Basdevant A, Oppert JM. One-year changes in energy expenditure and serum leptin following adjustable gastric banding in obese women. *Obes Surg* 2005;15:827–33.
- [24] Dargent J. Esophageal dilatation after laparoscopic adjustable gastric banding: definition and strategy. *Obes Surg* 2005;15:843–8.
- [25] De Luca M, de Werra C, Formato A, et al. Laparoscopic vs laparoscopic Lap-Band: 4-year results with early and intermediate complications. *Obes Surg* 2000;10:266–8.
- [26] De Waele B, Lauwers M, Van Nieuwenhove Y, Delvaux G. Out-patient laparoscopic gastric banding: initial experience. *Obes Surg* 2004;14:1108–10.
- [27] de Wit LT, Mathus-Vliegen L, Hey C, Rademaker B, Gouma DJ, Obertop H. Open versus laparoscopic adjustable silicone gastric banding: a prospective randomized trial for treatment of morbid obesity. *Ann Surg* 1999;230:800–5.
- [28] DeMaria EJ, Sugerman HJ, Meador JG, et al. High failure rate after laparoscopic adjustable silicone gastric banding for treatment of morbid obesity. *Ann Surg* 2001;233:809–18.
- [29] Dhafar KO. Initial experience with Swedish adjustable gastric band at Al-Noor hospital. *Obes Surg* 2003;13:918–20.

Appendix of Accepted Studies

- [1] Abu-Abeid S., Gavert N, Klausner JM, Szold A. Bariatric surgery in adolescence. *J Pediatr Surg* 2003;38:1379–82.
- [2] Abu-Abeid S, Keidar A, Gavert N, Blanc A, Szold A. The clinical spectrum of band erosion following laparoscopic adjustable silicone gastric banding for morbid obesity. *Surg Endosc* 2003;17:861–3.
- [3] Abu-Abeid S, Keidar A, Szold A. Resolution of chronic medical conditions after laparoscopic adjustable silicone gastric banding for the treatment of morbid obesity in the elderly. *Surg Endosc* 2001;15:132–4.
- [4] Al-Jiffry BO, Shaffer EA, Saccone GT, Downey P, Kow L, Toouli J. Changes in gallbladder motility and gallstone formation following laparoscopic gastric banding for morbid obesity. *Can J Gastroenterol* 2003;17:169–74.
- [5] Al-Momen A, El-Mogy I, Ibrahim A. Initial experience with Swedish adjustable gastric band at Saad Specialist Hospital, Al-Khobar, Saudi Arabia. *Obes Surg* 2005;15:506–9.
- [6] Albert S, Borovicka J, Thurnheer M, Lange J, Meyenberger C. Pre- and postoperative transaminase changes within the scope of gastric banding in morbid obesity. *Schweiz Rundsch Med Prax* 2001;90:1459–64.
- [7] Allen JW, Coleman MG, Fielding GA. Lessons learned from laparoscopic gastric banding for morbid obesity. *Am J Surg* 2001;182:10–4.
- [8] Angrisani L, Di Lorenzo N, Favretti F, Furbetta F, Iuppa A. Predictive value of initial body mass index for weight loss after 5 years of follow-up. *Surg Endosc* 2004;18:1524–7.
- [9] Belachew M, Belva PH, Desai C. Long-term results of laparoscopic adjustable gastric banding for the treatment of morbid obesity. *Obes Surg* 2002;12:564–8.

- [30] Doherty C, Maher JW, Heitshusen DS. Long-term data indicate a progressive loss in efficacy of adjustable silicone gastric banding for the surgical treatment of morbid obesity. *Surgery* 2002;132:724–7.
- [31] Dukhno O, Ovnat A, Levy I. Our experience with 250 laparoscopic adjustable silicone gastric bandings. *Surg Endosc* 2003;17:601–2.
- [32] Elakkary E, Elhorr A, Aziz F, Gazayerli MM, Silva YJ. Do support groups play a role in weight loss after laparoscopic adjustable gastric banding? *Obes Surg* 2006;16:331–4.
- [33] Elias B, Staudt JP, Van Vyne E. The technical approach in banding to avoid pouch dilatation. *Obes Surg* 2001;11:311–4.
- [34] Evans JD, Scott MH, Brown AS, Rogers J. Laparoscopic adjustable gastric banding for the treatment of morbid obesity. *Am J Surg* 2002;184:97–102.
- [35] Favretti F, Cadiere GB, Segato G, Himpens J, De Luca M, Busetto L. Laparoscopic banding: selection and technique in 830 patients. *Obes Surg* 2002;12:385–90.
- [36] Fielding GA, Duncombe JE. Clinical and radiological follow-up of laparoscopic adjustable gastric bands, 1998 and 2000: a comparison of two techniques. *Obes Surg* 2005;15:634–40.
- [37] Fielding GA, Rhodes M, Nathanson LK. Laparoscopic gastric banding for morbid obesity: surgical outcome in 335 cases. *Surg Endosc* 1999;13:550–4.
- [38] Fisher BL. Comparison of recovery time after open and laparoscopic gastric bypass and laparoscopic adjustable banding. *Obes Surg* 2004;14:67–72.
- [39] Forsell P, Hallerback B, Glise H, Hellers G. Complications following Swedish adjustable gastric banding: a long-term follow-up. *Obes Surg* 1999;9:11–6.
- [40] Fox SR, Fox KM, Srikanth MS, Rumbaut R. The Lap-Band system in a North American population. *Obes Surg* 2003;13:275–80.
- [41] Fransis K, Deweer F, Vanrykel JP, Aelvoet C. Experience with laparoscopic adjustable gastric banding (Lap-Band system) up to 8 years. *Acta Chir Belg* 2005;105:69–73.
- [42] Freys SM, Tigges H, Heimbucher J, Fuchs KH, Fein M, Thiede A. Quality of life following laparoscopic gastric banding in patients with morbid obesity. *J Gastrointest Surg* 2001;5:401–7.
- [43] Frigg A, Peterli R, Peters T, Ackermann C, Tondelli P. Reduction in co-morbidities 4 years after laparoscopic adjustable gastric banding. *Obes Surg* 2004;14:216–223.
- [44] Fruhbeck G, Rotellar F, Hernandez-Lizoain JL, et al. Fasting plasma ghrelin concentrations 6 months after gastric bypass are not determined by weight loss or changes in insulinemia. *Obes Surg* 2004;14:1208–15.
- [45] Furbetta F, Coli E. Codification of techniques for reoperation after Lap-Band. *Obes Surg* 2003;13:289–93.
- [46] Garrapa GG, Canibus P, Gatti C, et al. Changes in body composition and insulin sensitivity in severely obese subjects after laparoscopic adjustable silicone gastric banding (LASGB). *Med Sci Monit* 2005;11:CR522–8.
- [47] Gazzaruso C, Giordanetti S, La Manna A. Weight loss after Swedish adjustable gastric banding: relationships to insulin resistance and metabolic syndrome. *Obes Surg* 2002;12:841–5.
- [48] Greenslade J, Kow L, Tooouli J. Surgical management of obesity using a soft adjustable gastric band. *ANZ J Surg* 2004;74:195–9.
- [49] Greenstein RJ, Nissan A, Jaffin B. Esophageal anatomy and function in laparoscopic gastric restrictive bariatric surgery: implications for patient selection. *Obes Surg* 1998;8:199–206.
- [50] Hallerback B, Glise H, Johansson B, Johnson E. Laparoscopic surgery for morbid obesity. *Eur J Surg Suppl* 1998;582:128–31.
- [51] Hanusch-Enserer U, Hermann KM, Cauza E, Spak M, Mahr B, Dunky A. Effect of gastric banding on amino terminal pro-brain natriuretic peptide in the morbidly obese. *Obes Res* 2003;11:695–8.
- [52] Hell E, Miller KA, Moorehead MK, Norman S. Evaluation of health status and quality of life after bariatric surgery: comparison of standard Roux-en-Y gastric bypass, vertical banded gastroplasty, and laparoscopic adjustable silicone gastric banding. *Obes Surg* 2000;10:214–9.
- [53] Holeczy P, Novak P, Kralova A. 30% complications with adjustable gastric banding: what did we do wrong?. *Obes Surg* 2001;11:748–51.
- [54] Holloway JA, Forney GA, Gould DE. The Lap-Band is an effective tool for weight loss even in the United States. *Am J Surg* 2004;188:659–62.
- [55] Horschner R, Tuinebreijer MW, Kelder PH. Quality-of-life assessment of morbidly obese patients who have undergone a Lap-Band operation: 2-year follow-up study. Is the MOS SF-36 a useful instrument to measure quality of life in morbidly obese patients? *Obes Surg* 2001;11:212–8.
- [56] Horschner R, Tuinebreijer W, Kelder H. Eating patterns in morbidly obese patients before and after a gastric restrictive operation. *Obes Surg* 2002;12:108–12.
- [57] Iordache N, Vizetueu R, Iorgulescu A, Zmeu B, Iordache M. Laparoscopic adjustable gastric-banding treatment for morbid obesity: our first year experience. *Chirurgia (Bucur)* 2003;98:135–42.
- [58] Jan JC, Hong D, Pereira N, Patterson EJ. Laparoscopic adjustable gastric banding versus laparoscopic gastric bypass for morbid obesity: a single-institution comparison study of early results. *J Gastrointest Surg* 2005;9:30–41.
- [59] Jenkins JT, Modak P, Galloway DJ. Prospective study of laparoscopic adjustable gastric banding in the west of Scotland. *Scott Med J* 2006;51:37–41.
- [60] Kim TH, Daud A, Ude AO, et al. Early U.S. outcomes of laparoscopic gastric bypass versus laparoscopic adjustable silicone gastric banding for morbid obesity. *Surg Endosc* 2006;20:202–9.
- [61] Kirchmayr W, Klaus A, Muhlmann G, Mittermair R, Bonatti H, Aigner F. Adjustable gastric banding: assessment of safety and efficacy of bolus-filling during follow-up. *Obes Surg* 2004;14:387–91.
- [62] Korenkov M, Kneist W, Heintz A, Junginger T. Laparoscopic gastric banding as a universal method for the treatment of patients with morbid obesity. *Obes Surg* 2004;14:1123–7.
- [63] Kormanova K, Fried M, Hainer V, Kunesova M. Is laparoscopic adjustable gastric banding a day surgery procedure? *Obes Surg* 2004;14:1237–40.
- [64] Kothari SN, DeMaria EJ, Sugerman HJ, Kellum JM, Meador J, Wolfe L. Lap-Band failures: conversion to gastric bypass and their preliminary outcomes. *Surgery* 2002;131:625–9.
- [65] Laimer M, Kaser S, Kranebitter M, et al. Effect of pronounced weight loss on the nontraditional cardiovascular risk marker matrix metalloproteinase-9 in middle-aged morbidly obese women. *Int J Obes (Lond)* 2005;29:498–501.
- [66] Lap-Band PMA. The Lap-Band adjustable gastric banding system summary of safety and effectiveness data. Available at: <http://www.fda.gov/cdrh/pdf/P000008b.pdf>.
- [67] Larsen JF, Kroustrup JP. Laparoscopic adjustable gastric banding for the treatment of morbid obesity: six years' experiences. *Ugeskr Laeger* 2005;167:1946–9.
- [68] Larsen JK, Geenen R, Maas C, et al. Personality as a predictor of weight loss maintenance after surgery for morbid obesity. *Obes Res* 2004;12:1828–34.
- [69] Leonetti F, Silecchia G, Iacobellis G, Ribaudo MC, Zappaterreno A, Tiberti C. Different plasma ghrelin levels after laparoscopic gastric bypass and adjustable gastric banding in morbid obese subjects. *J Clin Endocrinol Metab* 2003;88:4227–31.
- [70] Mariani LM, Fusco A, Turriziani M, et al. Transient increase of plasma ghrelin after laparoscopic adjustable gastric banding in morbid obesity. *Horm Metab Res* 2005;37:242–5.
- [71] Martikainen T, Pirinen E, Alhava E, et al. Long-term results, late complications and quality of life in a series of adjustable gastric banding. *Obes Surg* 2004;14:648–654.

- [72] Meyer L, Rohr S, Becker J, et al. Retrospective study of laparoscopic adjustable silicone gastric banding for the treatment of morbid obesity: results and complications in 127 patients. *Diabetes Metab* 2004;30:53–60.
- [73] Miller K, Hell E. Laparoscopic adjustable gastric banding: a prospective 4-year follow-up study. *Obes Surg* 1999;9:183–7.
- [74] Mittermair RP, Weiss H, Nehoda H, Kirchmayr W, Aigner F. Laparoscopic Swedish adjustable gastric banding: 6-year follow-up and comparison to other laparoscopic bariatric procedures. *Obes Surg* 2003;13:412–7.
- [75] Mognol P, Chosidow D, Marmuse JP. Laparoscopic gastric bypass versus laparoscopic adjustable gastric banding in the super-obese: a comparative study of 290 patients. *Obes Surg* 2005;15:76–81.
- [76] Morino M, Toppino M, Bonnet G, del Genio G. Laparoscopic adjustable silicone gastric banding versus vertical banded gastroplasty in morbidly obese patients: a prospective randomized controlled clinical trial. *Ann Surg* 2003;238:835–41.
- [77] Moser F, Horgan S. Robotically assisted bariatric surgery. *Am J Surg* 2004;188:38S–44S.
- [78] Mouiel J. Morbid obesity, public health problem: Laparoscopic surgical treatment. *Bull Acad Natl Med* 2001;185:1475–90.
- [79] Nicolai A, Ippoliti C, Petrelli MD. Laparoscopic adjustable gastric banding: essential role of psychological support. *Obes Surg* 2002;12:857–63.
- [80] Nilsell K, Thorne A, Sjostedt S, Apelman J, Pettersson N. Prospective randomised comparison of adjustable gastric banding and vertical banded gastroplasty for morbid obesity. *Eur J Surg* 2001;167:504–9.
- [81] Nini E, Slim K, Scesa JL, Chipponi J. Evaluation of laparoscopic bariatric surgery using the BAROS score. *Ann Chir* 2002;127:107–14.
- [82] Neville E, Dams A, Vlasselaers J. Lap-Band erosion: incidence and treatment. *Obes Surg* 2001;11:744–7.
- [83] Nowara HA. Egyptian experience in laparoscopic adjustable gastric banding (technique, complications, and intermediate results). *Obes Surg* 2001;11:70–5.
- [84] O'Brien PE, Dixon JB. Laparoscopic adjustable gastric banding in the treatment of morbid obesity. *Arch Surg* 2003;138:376–82.
- [85] Paganelli M, Giacomelli M, Librenti MC, Pontiroli AE, Ferla G. Thirty months experience with laparoscopic adjustable gastric banding. *Obes Surg* 2000;10:269–71.
- [86] Perego L, Pizzocri P, Corradi D, et al. Circulating leptin correlates with left ventricular mass in morbid (grade III) obesity before and after weight loss induced by bariatric surgery: a potential role for leptin in mediating human left ventricular hypertrophy. *J Clin Endocrinol Metab* 2005;90:4087–93.
- [87] Phillips ML, Lewis MC, Chew V, et al. The early effects of weight loss surgery on regional adiposity. *Obes Surg* 2005;15:1449–55.
- [88] Ponce J, Paynter S, Fromm R. Laparoscopic adjustable gastric banding: 1,014 consecutive cases. *J Am Coll Surg* 2005;201:529–35.
- [89] Ponson AE, Janssen IM, Klinkenbijl JH. Laparoscopic adjustable gastric banding: a prospective comparison of two commonly used bands. *Obes Surg* 2002;12:579–82.
- [90] Ram E, Vishne T, Maayan R. The relationship between BMI, plasma leptin, insulin and proinsulin before and after laparoscopic adjustable gastric banding. *Obes Surg* 2005;15:1456–62.
- [91] Regusci L, Groebli Y, Meyer JL, Walder J, Margalith D, Schneider R. Gastroscopic removal of an adjustable gastric band after partial intragastric migration. *Obes Surg* 2003;13:281–4.
- [92] Ren CJ, Weiner M, Allen JW. Favorable early results of gastric banding for morbid obesity. *Surg Endosc* 2004;18:543–546.
- [93] Rosenthal RJ, Szomstein S, Kennedy CI, Soto FC, Zundel N. Laparoscopic surgery for morbid obesity: 1,001 consecutive bariatric operations performed at the Bariatric Institute, Cleveland Clinic Florida. *Obes Surg* 2006;16:119–24.
- [94] Rubenstein RB. Laparoscopic adjustable gastric banding at a US center with up to 3-year follow-up. *Obes Surg* 2002;12:380–4.
- [95] Rubin M, Spivak H. Prospective study of 250 patients undergoing laparoscopic gastric banding using the two-step technique. *Surg Endosc* 2003;17:857–60.
- [96] Sannen I, Himpens J, Leman G. Causes of dissatisfaction in some patients after adjustable gastric banding. *Obes Surg* 2001;11:605–8.
- [97] Sarker S, Myers J, Serot J, Shayani V. Three-year follow-up weight loss results for patients undergoing laparoscopic adjustable gastric banding at a major university medical center: does the weight loss persist? *Am J Surg* 2006;191:372–6.
- [98] Savastano S, Belfiore A, Guida B, et al. Role of dehydroepiandrosterone sulfate levels on body composition after laparoscopic adjustable gastric banding in pre-menopausal morbidly obese women. *J Endocrinol Invest* 2005;28:509–15.
- [99] Semple CW, Chehata A, Wilkinson S, Wertheimer MA. Laparoscopic adjustable gastric banding: initial Tasmanian experience. *ANZ J Surg* 2003;73:594–6.
- [100] Shapiro K, Patel S, Abdo Z, Ferzli G. Laparoscopic adjustable gastric banding: is there a learning curve? *Surg Endosc* 2004;18:48–50.
- [101] Silecchia G, Restuccia A, Elmore U, Polito D, Perrotta N, Genco A. Laparoscopic adjustable silicone gastric banding: prospective evaluation of intragastric migration of the Lap-Band. *Surg Laparosc Endosc Percutan Tech* 2001;11:229–34.
- [102] Silva AS, Cardoso H, Nogueira C, Santos J, Vilaca H. Treatment of morbid obesity with adjustable gastric band: preliminary report. *Obes Surg* 1999;9:194–7.
- [103] Skull AJ, Slater GH, Duncombe JE, Fielding GA. Laparoscopic adjustable banding in pregnancy: safety, patient tolerance, and effect on obesity-related pregnancy outcomes. *Obes Surg* 2004;14:230–235.
- [104] Spivak H, Anwar F, Burton S, Guerrero C, Onn A. The Lap-Band system in the United States. *Surg Endosc* 2004;18:198–202.
- [105] Spivak H, Hewitt MF, Onn A, Half EE. Weight loss and improvement of obesity-related illness in 500 U.S. patients following laparoscopic adjustable gastric banding procedure. *Am J Surg* 2005;189:27–32.
- [106] Steffen R, Biertho L, Ricklin T, Piec G, Horber FF. Laparoscopic Swedish adjustable gastric banding: a five-year prospective study. *Obes Surg* 2003;13:404–11.
- [107] Stieger R, Thurnheer M, Lange J. Morbid obesity: 130 consecutive patients with laparoscopic gastric banding. *Schweiz Med Wochenschr* 1998;128:1239–46.
- [108] Susmallian S, Ezri T, Charuzi I. Laparoscopic repair of access port site hernia after Lap-Band system implantation. *Obes Surg* 2002;12:682–4.
- [109] Suter M, Giusti V, Worreth M, Heraief E, Calmes JM. Laparoscopic gastric banding: a prospective, randomized study comparing the Lap-Band and the SAGB: early results. *Ann Surg* 2005;241:55–62.
- [110] Taskin M, Zengin K, Apaydin BB, Unal E. Laparoscopy in Turkish bariatric surgery: initial experience. *Obes Surg* 2000;10:263–5.
- [111] Taskin M, Zengin K, Unal E, Sakoglu N. Conversion of failed vertical banded gastroplasty to open adjustable gastric banding. *Obes Surg* 2001;11:731–4.
- [112] Thomusch O, Keck T, Dobschutz EV, Wagner C, Ruckauer KD, Hopt UT. Risk factors for the intermediate outcome of morbid obesity after laparoscopically placed adjustable gastric banding. *Am J Surg* 2005;189:214–8.
- [113] Thorne A, Lonnqvist F, Apelman J, Hellers G, Arner P. A pilot study of long-term effects of a novel obesity treatment: omentectomy in connection with adjustable gastric banding. *Int J Obes Relat Metab Disord* 2002;26:193–9.
- [114] Ti TK. Singapore experience in obesity surgery. *Obes Surg* 2004;14:1103–7.
- [115] Toppino M, Morino M, Bonnet G, Nigra I, Siliquini R. Laparoscopic surgery for morbid obesity: preliminary results from SICE registry

- (Italian Society of Endoscopic and Minimally Invasive Surgery). *Obes Surg* 1999;9:62–5.
- [116] Tweddle EA, Woods S, Blamey S. Laparoscopic gastric banding: safe and modestly successful. *ANZ J Surg* 2004;74:191–4.
- [117] van Dielen FM, Soeters PB, de Brauw LM, Greve JW. Laparoscopic adjustable gastric banding versus open vertical banded gastroplasty: a prospective randomized trial. *Obes Surg* 2005;15:1292–8.
- [118] Vassallo C, Andreoli M, La Manna A, Turpini C. 60 reoperations on 890 patients after gastric restrictive surgery. *Obes Surg* 2001;11:752–6.
- [119] Vertruyen M, Paul G. 11-cm Lap-Band system placement after history of intragastric migration. *Obes Surg* 2003;13:435–8.
- [120] Victorzon M, Tolonen P. Intermediate results following laparoscopic adjustable gastric banding for morbid obesity. *Dig Surg* 2002;19:354–7.
- [121] Wasserberg N, Nudelman I, Fuko Z, Rubin M. Laparoscopic repositioning of Heliogast gastric band after anterior slippage. *Obes Surg* 2003;13:780–3.
- [122] Watkins BM, Montgomery KF, Ahroni JH. Laparoscopic adjustable gastric banding: early experience in 400 consecutive patients in the USA. *Obes Surg* 2005;15:82–7.
- [123] Weber M, Muller MK, Bucher T, et al. Laparoscopic gastric bypass is superior to laparoscopic gastric banding for treatment of morbid obesity. *Ann Surg* 2004;240:975–83.
- [124] Weiner R, Blanco-Engert R, Weiner S, Matkowitz R, Schaefer L, Pomhoff I. Outcome after laparoscopic adjustable gastric banding: 8 years experience. *Obes Surg* 2003;13:427–34.
- [125] Wenger M, Piec G, Branson R, Potoczna N, Horber FF, Steffen R. Salvage of gastric restriction following staple-line dehiscence after vertical banded gastroplasty by insertion of an adjustable gastric band. *Obes Surg* 2005;15:216–22.
- [126] Wiesner W, Schlumpf R, Schob O, Hauser R, Kacel GM. Gastric pouch dilatation: complications after laparoscopic implantation of a silicone gastric band in pathologic obesity. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 1998;169:479–83.
- [127] Wylezol M, Pardela M, Gluck M, Puzio J, Czapla J. Swedish adjustable gastric band (SAGB) implanted laparoscopically in the treatment of morbid obesity: the first experience in Poland. *Med Sci Monit* 2000;6:441–5.
- [128] Zacharoulis D, Roy-Chadhury H, Dobbins B, et al. Laparoscopic adjustable gastric banding: surgical and radiological approach. *Obes Surg* 2002;12:280–4.
- [129] Zehetner J, Holzinger F, Triaca H, Klaiber C. A 6-year experience with the Swedish adjustable gastric band: prospective long-term audit of laparoscopic gastric banding. *Surg Endosc* 2005;19:21–8.